



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## MECHANICS.

173. Proposed by J. F. LAWRENCE, A. B., Professor of Mathematics, Oklahoma Agricultural College, Stillwater, Oklahoma.

A squirrel is in a cylindrical cage and oscillating with it about its axis which is horizontal. At the instant when he is at the highest point of the oscillation, he leaps to the opposite extremity of the diameter and arrives there at the same instant as the point at which he left. Determine his leap completely.

[No satisfactory solution has been received.]

174. Proposed by F. P. MATZ, Sc. D., Ph. D., Reading, Pa.

By what per cent. is the striking force of a hailstone increased in falling 1000 feet through a stratum of atmosphere moving uniformly eastward at the rate of 60 miles an hour?

Solution by the PROPOSER.

Let  $S_1$  = the *striking-force* of the hail-stone falling 1000 feet, and  $S_2 = mS_1$  = that caused by the horizontal motion of the atmosphere during the time required to fall 1000 feet; then the resultant striking-force is  $\sqrt{(1+m^2)}S_1$ . Assuming a mass of ice suitable for a hail-stone, we can easily calculate  $m$ .

The required percentage of increase in striking-force becomes

$$I = \left( \frac{\sqrt{(1+m^2)} - 1}{1} \right) \text{ of } 100\%.$$

175. Proposed by J. F. LAWRENCE, A. B., Professor of Mathematics, Oklahoma Agricultural College, Stillwater, Oklahoma.

A cylinder descends down a plane, the inclination of which to the horizon is  $\alpha$ , unwrapping a fine string fixed at the highest point of the plane. Find the angle through which the plane must be depressed in order that a sphere, descending under like circumstances, may experience the same acceleration.

Solution by G. W. GREENWOOD, M. A., Professor of Mathematics and Astronomy, McKendree College, Lebanon, Ill.

Let  $\omega$  be the angular velocity, and  $v$  the velocity parallel to the plane, after rolling a distance  $l$  from the highest point. Equating the *vis viva* of the body to twice the work done by gravity, we get, in the case of the cylinder,

$$\frac{1}{2}ma^2\omega^2 + mv^2 = 2mgl\sin\alpha,$$

where  $a$  is the radius of the cylinder.

$$\therefore 3v^2 = 4gl\sin\alpha, \text{ since } a\omega = v.$$

In the case of the sphere we get

$$\frac{2}{5}m'a'\omega'^2 + m'v'^2 = 2m'gl\sin(\alpha - \theta), \text{ i. e., } 7v'^2 = 10gl\sin(\alpha - \theta),$$

where  $\theta$  is the amount of depression of the plane.

$$\therefore 14\sin\alpha = 15\sin(\alpha - \theta), \text{ or } \theta = \alpha - \sin^{-1}\left(\frac{14}{15}\sin\alpha\right).$$